arXiv:0906.0654v1 [math.AG] 3 Jun 2009

On the birational geometry for irreducible symplectic 4-folds related to the Fano schemes of lines

Kotaro Kawatani*

Osaka University

1 Introduction

A compact Kähler manifold X is said to be an irreducible symplectic manifold when X is a simply connected and $H^0(X, \Omega_X^2)$ is generated by an everywhere non-degenerate holomorphic 2-form σ_X. We call such a holomorphic 2-form σ_X a symplectic form. Let Y be a smooth cubic 4-fold and $F(Y)$ the set of all lines contained in Y. $F(Y)$ is said to be the Fano schemes of lines. Beauville and Donagi [B-D] showed that $F(Y)$ is an irreducible symplectic 4-fold. Examples of irreducible symplectic manifolds are few. Most of them are constructed as the moduli spaces of stable sheaves on K3 surfaces or Abelian surfaces.

Assume that a finite group G acts on $F(Y)$. If G preserves a symplectic form $\sigma_{F(Y)}$ then there exists a symplectic form $\bar{\sigma}$ on the smooth locus of $X_G := F(Y)/G$. In general X_G has singular points. Let $\nu : \tilde{X}_G \to X_G$ be a resolution of X_G. The symplectic 2-form $\bar{\sigma}$ lifts to an everywhere non-degenerate 2-form on $\tilde{X}_G$ if and only if the resolution ν is crepant. If ν is crepant, then $\tilde{X}_G$ is also an irreducible symplectic 4-fold. The G-action on $F(Y)$ is said to be a good action when G preserves $\sigma_{F(Y)}$ and X_G has a crepant resolution. Can we find good actions on $F(Y)$? When it is possible, what is $\tilde{X}_G$?

In this paper, we assume that G is a subgroup of the automorphism group $PGL(5)$ of $\mathbb{P}^5$ and G preserves Y. We study two examples of good actions on $F(Y)$. In the first example, the resolution $\tilde{X}_G$ is birational to the generalized

---

*kawatani@cr.math.sci.osaka-u.ac.jp

Kummer variety $K^2(A)$ of an Abelian surface A. In the second one, $\tilde{X}_G$ is birational to the 2-points Hilbert scheme $\mathrm{Hilb}^2(S)$ of a K3 surface S. We have to investigate the G-action on symplectic forms on $F(Y)$. The Abel-Jacobi map defines a Hodge isomorphism of type $(-1,-1)$ from $H^4(Y,\mathbb{C})$ to $H^2(F(Y),\mathbb{C})$. Using the identification with $H^0(F(Y),\Omega^2_{F(Y)})$ and $H^1(Y,\Omega^3_Y)$, we can check whether G preserves the symplectic form or not (see Lemma 2.3).

In Section 3, we consider the first example, which was found by Namikawa ([Nam], 1.7.(iv)). Let Y be a smooth cubic 4-fold defined by

$$\{(z_0:\cdots:z_5)\in\mathbb{P}^5 | f(z_0,z_1,z_2)+g(z_3,z_4,z_5)=0\},$$

where f and g are homogeneous cubic polynomials. In addition, we consider two elliptic curves defined by $C:=\{(z_0:z_1:z_2)\in\mathbb{P}^2|f(z_0,z_1,z_2)=0\}$ and $D:=\{(z_3:z_4:z_5)\in\mathbb{P}^2|g(z_3,z_4,z_5)=0\}$. Assume that G is the cyclic group $\mathbb{Z}_3$ of order 3. The group G acts on Y in the following way :

$$\mathbb{Z}_3 \curvearrowright Y \ni (z_0:\cdots:z_5)\mapsto(z_0:z_1:z_2:\zeta z_3:\zeta z_4:\zeta z_5)$$

where $\zeta=\exp(2\pi\sqrt{-1}/3)$. $X_{\mathbb{Z}_3}$ has singularities of type A_2 along $C\times D$ and is smooth otherwise. So $X_{\mathbb{Z}_3}$ has a crepant resolution $\tilde{X}_{\mathbb{Z}_3}$ (Lemma 3.1). $\tilde{X}_{\mathbb{Z}_3}$ is birational to $K^2(C\times D)$ for the elliptic curves C and D (Proposition 3.2). There is a natural birational map $\psi:\tilde{X}_{\mathbb{Z}_3}\dashrightarrow K^2(C\times D)$ which is the Mukai flop on disjoint 18 copies of $\mathbb{P}^2$ (Theorem 3.5). Furthermore when C and D are not isogenus, we found 2 more symplectic birational models X_{I} and X_{II} other than $\tilde{X}_{\mathbb{Z}_3}$ and $K^2(C\times D)$ (Theorem 3.11).

In Section 4, we consider the second example. Let Y be a smooth cubic 4-fold defined by $\{(z_0:\cdots:z_5)\in\mathbb{P}^5|z_0^3+\cdots+z_5^3=0\}$. Put

$$M=\left\{A=\begin{pmatrix}a_1 & & \\ & \ddots & \\ & & a_6\end{pmatrix}\in PGL(5) | \#\{a_i=1\}=\#\{j|a_j=\zeta\}=3\right\}.$$

Let G be the finite group generated by M. Then G is isomorphic to $\mathbb{Z}_3^{\oplus 4}$ and the G-action on $F(Y)$ turns out to be good. Let E be the elliptic curve defined by $\{(z_0:z_1:z_2)\in\mathbb{P}^2|z_0^3+z_1^3+z_2^3=0\}$. Since E has a complex multiplication $\zeta \curvearrowright E\ni x\mapsto \zeta x$, the Abelian surface $E\times E$ has the $\mathbb{Z}_3$ action given by $\mathbb{Z}_3\curvearrowright E\times E\ni(x,y)\mapsto(\zeta x,\zeta^2 y)\in E\times E$. This $\mathbb{Z}_3$ action on $E\times E$ preserves a symplectic form on $E\times E$, and $(E\times E)/\mathbb{Z}_3$ has a crepant resolution $S\to(E\times E)/\mathbb{Z}_3$. In particular S is a K3 surface. We shall prove that $\tilde{X}_G$ is birational to $\mathrm{Hilb}^2(S)$ (Theorem 4.2).

Acknowledgment. The author would like to thank Professor Namikawa for his advice.

2 Preliminaries

In this section we prepare some facts needed later.

2.1 On irreducible symplectic manifolds

Let A be an Abelian surface and $\mathrm{Hilb}^{n+1}(A)$ the Hilbert scheme of $(n+1)$-points on A. We define

$$\mathrm{Sym}^{n+1}(A) := A^{n+1}/\mathfrak{S}_{n+1}$$

where $\mathfrak{S}_{n+1}$ is the $(n+1)$-th symmetric group. Let μ be the Hilbert-Chow morphism:

$$\mu : \mathrm{Hilb}^{n+1}(A) \to \mathrm{Sym}^{n+1}(A).$$

We define the map $\Sigma : \mathrm{Sym}^{n+1}(A) \to A$ by $\Sigma(\{x_1, \cdots x_{n+1}\}) := \sum_{i=1}^{n+1} x_i$ and consider the composite $\Sigma \circ \mu$. We define :

$$K^n(A) := (\Sigma \circ \mu)^{-1}(0).$$

We call $K^n(A)$ *the generalized Kummer variety of* A. $K^n(A)$ is an irreducible symplectic manifold ([Bea]). We also define $\bar{K}^n(A) := \Sigma^{-1}(0)$. $\bar{K}^n(A)$ is isomorphic to

$$\bar{K}^n(A) \cong \{(x_1, \cdots, x_{n+1}) \in A^{n+1} | x_1 + \cdots + x_{n+1} = 0\}/\mathfrak{S}_{n+1}.$$

For an elliptic curve C, we define

$$K^2(C) := \big\{\{p_1, p_2, p_3\} \in \mathrm{Hilb}^3(C) | p_1 + p_2 + p_3 = 0\big\}.$$

Notice that $K^2(C)$ is isomorphic to $\mathbb{P}^2$.

2.2 Fano schemes

We collect some facts for Fano schemes needed later. Let Y be a smooth cubic 4-fold. Define

$$F(Y) := \{l \subset Y | l \cong \mathbb{P}^1, \deg l = 1\}.$$

$F(Y)$ is known to be a smooth projective variety of dimension 4. Furthermore we have the following theorem.

Proposition 2.1 (Beauville-Donagi[B-D]). *The notation being as above, $F(Y)$ is an irreducible symplectic manifold. $F(Y)$ is deformation equivalent to the Hilbert scheme $\mathrm{Hilb}^2(S)$ of 2-point on a K3 surface S.*

The following relation between $H^4(Y, \mathbb{C})$ and $H^2(F(Y), \mathbb{C})$ is important to us.

Lemma 2.2 ([B-D])**.** *Let $\mathcal{L}$ be the universal family of lines*:

$$\mathcal{L} := \{(l, y) \in F(Y) \times Y | l \ni y\}.$$

Let p and q be the projections from $\mathcal{L}$ to $F(Y)$ and Y respectively. We define the Abel-Jacobi map α by

$$\alpha : H^4(Y, \mathbb{C}) \to H^2(F(Y), \mathbb{C}),\ \alpha(\omega) := p_* q^*(\omega).$$

Then α is a Hodge isomorphism of type $(-1, -1)$*.*

Let $(z_0 : \ldots : z_5)$ be the homogeneous coordinates of $\mathbb{P}^5$, and $f_Y(z_0, \ldots, z_5)$ the defining polynomial of Y. Let $Res : H^5(\mathbb{P}^5 - Y, \mathbb{C}) \to H^4(Y, \mathbb{C})$ be the residue map. By Lemma 2.2, we have $\alpha(H^1(Y, \Omega^3_Y)) = H^0(F(Y), \Omega^2_{F(Y)})$. Recall that $H^1(Y, \Omega^3_Y)$ is generated by $Res\frac{\Omega}{f_Y^2}$ ([Gri]), where $\Omega = \sum_{i=0}^5 (-1)^i z_i dz_0 \wedge \cdots \check{dz_i} \cdots \wedge dz_5$. From these arguments, we get the following commutative diagram:

$$\begin{tikzcd}
H^4(Y, \mathbb{C}) \arrow[r, "\cong", "\alpha"'] & H^2(F(Y), \mathbb{C}) \\
H^1(Y, \Omega^3_Y) \arrow[u, hook] \arrow[r] & H^0(F(Y), \Omega^2_{F(Y)}) \arrow[u, hook] \\
\mathbb{C}\langle Res\frac{\Omega}{f_Y^2}\rangle \arrow[u, equal] \arrow[r] & \mathbb{C}\langle \sigma_{F(Y)} \rangle. \arrow[u, equal]
\end{tikzcd}$$

2.3 Group actions on $F(Y)$

Let $PGL(5)$ be the automorphism group of $\mathbb{P}^5$, and G a finite subgroup of $PGL(5)$. We assume that G preserves Y. We need a criterion for the action preserving the symplectic form.

Let us consider the induced action $G \curvearrowright F(Y)$. Since the Abel-Jacobi map α is G-equivariant, the induced action $G \curvearrowright F(Y)$ preserves the symplectic form $\sigma_{F(Y)}$ if and only if G preserves $Res\frac{\Omega}{f_Y^2}$. So, we get the following lemma.

Lemma 2.3. *Assume that G preserves Y. Then G preserves the symplectic form $\sigma_{F(Y)}$ on $F(Y)$ if and only if G preserves $Res\frac{\Omega}{f_Y^2}$*

3 First example

We first recall results of [Nam]. We define a smooth cubic 4-fold Y by

$$Y := \{(z_0 : \cdots : z_5) \in \mathbb{P}^5 | f(z_0, z_1, z_2) + g(z_3, z_4, z_5) = 0\}.$$

Define two elliptic curves by

$$C := \{(z_0 : z_1 : z_2) \in \mathbb{P}^2 | f(z_0, z_1, z_2) = 0\}$$

and

$$D := \{(z_3 : z_4 : z_5) \in \mathbb{P}^2 | g(z_3, z_4, z_5) = 0\}.$$

We put

$$P_C = \{(z_0 : \cdots : z_5) \in \mathbb{P}^5 | z_3 = z_4 = z_5 = 0\},$$

and

$$P_D = \{(z_0 : \cdots : z_5) \in \mathbb{P}^5 | z_0 = z_1 = z_2 = 0\}.$$

Notice that $C \subset P_C$ and $D \subset P_D$. Assume that $G \subset PGL(5)$ is the cyclic group $\mathbb{Z}_3$ of order 3 generated by

$$\tau = \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ & & & \zeta & & \\ & & & & \zeta & \\ & & & & & \zeta \end{pmatrix},$$

where $\zeta = \exp(2\pi\sqrt{-1}/3)$.

Let us consider the natural group action :

$$\mathbb{Z}_3 \curvearrowright \mathbb{P}^5 \ni (z_0 : z_1 : z_2 : z_3 : z_4 : z_5) \mapsto (z_0 : z_1 : z_2 : \zeta z_3 : \zeta z_4 : \zeta z_5).$$

Since $\mathbb{Z}_3$ preserves Y, the action induces the $\mathbb{Z}_3$-action on $F(Y)$.

Lemma 3.1. *The induced action $\mathbb{Z}_3 \curvearrowright F(Y)$ preserves the symplectic form $\sigma_{F(Y)}$, and the singular locus of $X_{\mathbb{Z}_3} := F(Y)/\mathbb{Z}_3$ is isomorphic to $C \times D$. The singularity is A_2. In particular $X_{\mathbb{Z}_3}$ has a crepant resolution $\nu : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$.*

Proof. By Lemma 2.3, the induced action $\mathbb{Z}_3 \curvearrowright F(Y)$ preserves the symplectic form $\sigma_{F(Y)}$. The singular locus of $X_{\mathbb{Z}_3}$ is isomorphic to the fixed locus $\mathrm{Fix}_{\mathbb{Z}_3}(F(Y))$ of the $\mathbb{Z}_3$ action on $F(Y)$. We remark that $\mathrm{Fix}_{\mathbb{Z}_3}(Y)$ is the disjoint union $C \cup D$. If a line l is in $\mathrm{Fix}_{\mathbb{Z}_3}(F(Y))$, then $\mathbb{Z}_3$ acts on the line l. l has two fixed points by the $\mathbb{Z}_3$ action. So $l \in F(Y)$ is in $\mathrm{Fix}_{\mathbb{Z}_3}(F(Y))$ if and only if l passes through a point $p \in C$ and $q \in D$. Hence $\mathrm{Fix}_{\mathbb{Z}_3}(F(Y))$ is isomorphic to $C \times D$.


P_C
$C = \{f(z_0, z_1, z_2) = 0\}$
p
l
$\mathrm{Fix}_{\mathbb{Z}_3}(F(Y)) \cong \{l = \langle p, q\rangle | p \in C,\ q \in D\}$
$D = \{g(z_3, z_4, z_5) = 0\}$
q
P_D

Since $\mathbb{Z}_3$ preserves $\sigma_{F(Y)}$, $X_{\mathbb{Z}_3}$ has A_2 singularities along $C \times D$. So, $X_{\mathbb{Z}_3}$ has a crepant resolution $\nu : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$. ☐

Proposition 3.2 ([Nam] 1.7.(iv))**.** *The notation being as above, $\tilde{X}_{\mathbb{Z}_3}$ is birational to $K^2(C \times D)$.*

Proof. We first define a rational map $\psi : \tilde{X}_{\mathbb{Z}_3} \dashrightarrow K^2(C \times D)$. Let l be in $F(Y)$, and $[l]$ the orbit of l by the $\mathbb{Z}_3$ action : $[l] = \{l, \tau(l), \tau^2(l)\}$. Let W_l be the linear space spanned by $l, \tau(l)$ and $\tau^2(l)$. If we choose $l \in F(Y)$ in general, then W_l is isomorphic to $\mathbb{P}^3$ and both $P_C \cap W_l$ and $P_D \cap W_l$ are lines. Furthermore we may assume that the cubic surface $S := W_l \cap Y$ is smooth. There are 27 lines in S [Har]. Notice that three lines $l, \tau(l)$ and $\tau^2(l)$ are contained in S.

We will prove follwoings.

1. There are three lines m_1, m_2 and m_3 such that m_i does not meet m_j $(i \neq j)$, and each m_i meets l at one point (respectively $\tau(l), \tau^2(l)$). See also Figure 1.

2. The set of three lines is unique, and

3. each m_i $(i = 1, 2, 3)$ is fixed by the $\mathbb{Z}_3$ action.

Indeed the cubic surface S is the blow up of $\mathbb{P}^2$ along 6 points $a_1, \ldots, a_6$. Let E_i be the exceptional (-1)-curve over a_i, C_i the proper transform of the conic passing through 5 points except a_i, and L_{ij} the proper transform of line passing through a_i and a_j. We may assume that $E_1 = l$, $E_2 = \tau(l)$ and $E_3 = \tau^2(l)$. By the configuration of lines, each m_i is given by $m_1 = C_4, m_2 = C_5$ and $m_3 = C_6$. There are exactly six lines which does not intersect $\tau^k(l)$ $(k = 0, 1, 2)$. These six lines are given by $L_{45}, L_{46}, L_{56}, E_4, E_5$ and E_6. Since $\mathbb{Z}_3$ acts on the set $\{E_1, E_2, E_3\}$, $\mathbb{Z}_3$ acts on the set $\{L_{45}, L_{46}, L_{56}, E_4, E_5, E_6\}$. Since the order of $\mathbb{Z}_3$ is 3, $\mathbb{Z}_3$ acts on the two set $\{L_{45}, L_{46}, L_{56}\}$ and $\{E_4, E_5, E_6\}$.

If E_4, E_5 and E_6 are not fixed by $\mathbb{Z}_3$ then all lines in S move by the $\mathbb{Z}_3$ action. Since there are 9 lines in S fixed by the $\mathbb{Z}_3$ action, three lines E_4, E_5 and E_6 are fixed by the $\mathbb{Z}_3$ action. Hence three lines m_1, m_2 and m_3 are fixed by the $\mathbb{Z}_3$ action. In particular m_i $(i = 1, 2, 3)$ meet C (resp D) at one point.

Put $p_i = m_i \cap C$ and $q_i = m_i \cap D$. Since three points $\{p_1, p_2, p_3\}$(resp. $\{q_1, q_2, q_3\}$) are in the line $P_C \cap W_l$, the sum $p_1 + p_2 + p_3$ is $0 \in C$ (resp. $q_1 + q_2 + q_3 = 0 \in D$). So we get an element $\{(p_i, q_i)\}_{i=1}^3$ of $K^2(C \times D)$. We define ψ as:



Figure 1: Picture of ψ and φ

$$\psi : \tilde{X}_{\mathbb{Z}_3} \dashrightarrow K^2(C \times D), \psi([l]) = \{(p_i, q_i)\}_{i=1}^3.$$

We next define a rational map $\varphi : K^2(C \times D) \dashrightarrow \tilde{X}_{\mathbb{Z}_3}$. Let $\{(p_i, q_i)\}_{i=1}^3$ be in $K^2(C \times D)$. We may assume that $p_i \neq p_j$ and $q_i \neq q_j$ $(i \neq j)$. Let $m_i (i = 1, 2, 3)$ be the line passing through p_i and q_i. Let M be the linear space spanned by m_1, m_2 and m_3. Then M is isomorphic to $\mathbb{P}^3$. Let S be the cubic surface $M \cap Y$. We remark that S is smooth. By the configuration of 27 lines, there are exactly three lines l_1, l_2 and l_3 such that $l_i \cap l_j = \emptyset$ $(i \neq j)$, and each l_i meets m_j $(i, j = 1, 2, 3)$ at exactly one point. We notice that m_1, m_2 and m_3 are fixed by the $\mathbb{Z}_3$ action on S. By the uniqueness of the set of three lines, we have

$$\mathbb{Z}_3 \curvearrowright \{l_1, l_2, l_3\} \mapsto \{l_1, l_2, l_3\}.$$

Since each l_i is not fixed by the $\mathbb{Z}_3$ action, we may assume that $\tau(l_1) = l_2$.

By the uniqueness of the set of three lines, both φ and ψ are birational maps. In particular, $\varphi^{-1} = \psi$. □

3.1 On the indeterminacy of the birational maps

In Theorem 3.5 we will prove that φ and ψ is given as the Mukai flop on 18 disjoint copies of $\mathbb{P}^2$. The key lemma of the proof is Lemma 3.3. In the proof of Lemma 3.3, we give the geometric description of the exceptional divisor of $\nu : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$ by using G-Hilbert scheme. Let X be a smooth projective variety over $\mathbb{C}$. We assume that a finite group G acts on X. Then we define G-Hilb(X) as

$$G\text{-Hilb}(X) := \{Z \in \text{Hilb}^{\#G}(X) | \mathcal{O}_Z \cong \mathbb{C}[G] \text{ (as } G \text{ modules)}\}.$$

and we call it *G-Hilbert scheme of X*. Let z be in $\tilde{X}_{\mathbb{Z}_3}$ and we assume that z is not in the exceptional locus Exc(ν) of ν. Then z is the orbit of $l \in F(Y)$ by the $\mathbb{Z}_3$ action. Hence z is identified with $Z \in \mathbb{Z}_3$-Hilb$(F(Y))$. Assume that z is in Exc(ν). On a neighborhood of Fix$(\mathbb{Z}_3)$, the $\mathbb{Z}_3$ action can be identified with

$$\mathbb{Z}_3 \curvearrowright \text{Spec}\mathbb{C}[x_1, x_2, x_3, x_4], \ (x_1, x_2, x_3, x_4) \mapsto (x_1, x_2, \zeta x_3, \zeta^2 x_4).$$

By using the G-graph, we see that $\mathbb{Z}_3$-Hilb(Spec$\mathbb{C}[x_1, x_2, x_3, x_4]$) is isomorphic to $\mathbb{C}^2 \times \mathbb{Z}_3$-Hilb$(\mathbb{C}^2)$ (See [Nak]). So $\mathbb{Z}_3$-Hilb(Spec$\mathbb{C}[x_1, x_2, x_3, x_4]$) is a crepant resolution of $\mathbb{C}^4/\mathbb{Z}_3$. Since this crepant resolution is unique, the element z of $\tilde{X}_{\mathbb{Z}_3}$ can be identified with the element Z of $\mathbb{Z}_3$-Hilb$(F(Y))$. By pulling back the universal family $\mathcal{L}$ of $F(Y)$, we get the family of lines $\mathcal{L}_Z$ over Z.

$$\begin{tikzcd}
\mathcal{L}_Z \arrow[r] \arrow[d] & \mathcal{L} \subset F(Y) \times \mathbb{P}^5 \arrow[r] \arrow[d] & \mathbb{P}^5 \\
Z \arrow[r] & F(Y) &
\end{tikzcd}$$

The support of $\mathcal{L}_Z$ is $l \cup \tau(l) \cup \tau^2(l)$, where $l \in F(Y)$. Here we put $[l] := l \cup \tau(l) \cup \tau^2(l)$. When l is not in the fixed locus of the $\mathbb{Z}_3$ action, $\mathcal{L}_Z$ is uniquely determined by $[l]$. If l is in the fixed locus of the $\mathbb{Z}_3$ action, then Z is not determined uniquely by $[l]$.

Let p_2 be the second projection $p_2 : F(Y) \times \mathbb{P}^5 \to \mathbb{P}^5$. For each Z, take the smallest linear subspace L contained in $\mathbb{P}^5$ such that $\mathcal{L}_Z \subset p_2^{-1}(L)$. We call this L the linear space spanned by $\mathcal{L}_Z$ (or simply Z). In a generic case, Z spans $\mathbb{P}^3$. In some special cases Z spans $\mathbb{P}^2$.

Lemma 3.3. *Assume that the line $l \in F(Y)$ is in the fixed locus of the $\mathbb{Z}_3$ action, that is, l passes through $p \in C$ and $q \in D$. We only consider $Z \in \tilde{X}_{\mathbb{Z}_3}$ such that the support of $\mathcal{L}_Z$ is l. Then Z is parametrized by the tree $\mathcal{I} \cup \mathcal{J}$ of two projective lines $\mathcal{I}$ and $\mathcal{J}$.*

1. Assume that neither p nor q are 3-torsion points. Then $\mathcal{L}_Z$ spans $\mathbb{P}^3$, for each Z.

2. Assume that both p and q are 3-torsion points. There exists exactly one point on each $\mathbb{P}^1$ such that $\mathcal{L}_Z$ spans $\mathbb{P}^2$. These two points are not in the intersection $\mathcal{I} \cap \mathcal{J}$. For other points, $\mathcal{L}_Z$ spans $\mathbb{P}^3$.

3. Assume that p is a 3-torsion point and q is not a 3-torsion point. There exists exactly one point on the tree such that $\mathcal{L}_Z$ spans $\mathbb{P}^2$. Moreover this point is not in the intersection $\mathcal{I} \cap \mathcal{J}$. Otherwise $\mathcal{L}_Z$ spans $\mathbb{P}^3$.

When q is a 3-torsion point and p is not a 3-torsion point, we have a similar situation to *3*.

Proof. Let $Z \in \tilde{X}_{\mathbb{Z}_3}$ be a closed subscheme of length 3 such that the support of $\mathcal{L}_Z$ is the line passing through $p \in C$ and $q \in D$. Let $(x : y : z : u : v : w)$ be the homogeneous coordinates of $\mathbb{P}^5$, consider the cubic

$$Y := \{(x : \cdots : w) \in \mathbb{P}^5 | f(x, y, z) + g(u, v, w) = 0\}.$$

Let C and D be elliptic curves defined by

$$C := \{(x : y : z) \in \mathbb{P}^2 | f(x, y, z) = 0\},\ D := \{(u : v : w) \in \mathbb{P}^2 | g(u, v, w) = 0\}.$$

We put

$$P_C := \{(x : \cdots : w) \in \mathbb{P}^5 | u = v = w = 0\},$$

and

$$P_D := \{(x : \cdots : w) \in \mathbb{P}^5 | x = y = z = 0\}.$$

1. By changing coordinates, we may assume that

$$p = (0 : 0 : 1),\ f(x, y, z) = x(x - z)(x - \alpha z) - y^2 z,\ \alpha \in \mathbb{C} - \{0, 1\}$$

$$q = (0 : 0 : 1),\ g(u, v, w) = u(u - w)(u - \beta w) - v^2 w,\ \ \beta \in \mathbb{C} - \{0, 1\}.$$

So the line l is given by:

$$l = (0 : 0 : \lambda : 0 : 0 : \mu) \text{ where } (\lambda : \mu) \in \mathbb{P}^1.$$

Let $G(2, 6)$ be the Grassmann manifold parameterizing 2-dimensional subspace of $\mathbb{C}^6$. We will give coordinates system around l by using the local coordinates of $G(2, 6)$. We define two hyperplanes

$$\{z = 0\} := \{(x : y : \cdots : w) \in \mathbb{P}^5 | z = 0\},$$
$$\{w = 0\} := \{(x : y : \cdots : w) \in \mathbb{P}^5 | w = 0\}.$$

Then the intersection $\{w = 0\} \cap l$ is $p = (0 : 0 : 1 : 0 : 0 : 0)$ and the intersection $\{z = 0\} \cap l$ is $q = (0 : 0 : 0 : 0 : 0 : 1)$. Let (u_0, u_1, u_3, u_4) be affine coordinates of the hyperplane $\{w = 0\}$ around p :

$$(u_0 : u_1 : 1 : u_3 : u_4 : 0) \in \mathbb{P}^4.$$

Let (v_0, v_1, v_3, v_4) be affine coordinates of the hyperplane $\{z = 0\}$ around q :

$$(v_0 : v_1 : 0 : v_3 : v_4 : 1) \in \mathbb{P}^4.$$

Let U_l be an open neighborhood of l in $G(2,6)$. The local coordinates of U_l is given by $(u_0, u_1, u_3, u_4, v_0, v_1, v_3, v_4)$. Indeed, for each $(u_0, \cdots, v_4) \in U_l$, the corresponding line $\tilde{l}$ is given by

$$(u_0, \cdots, v_4) \leftrightarrow \tilde{l} := \lambda(u_0 : u_1 : 1 : u_3 : u_4 : 0) + \mu(v_0 : v_1 : 0 : v_3 : v_4 : 1)$$

The line $\tilde{l} \in G(2,6)$ is in $F(Y)$ if and only if $\tilde{l}$ is contained in Y. Therefore we have

$$(f+g)(\tilde{l}) = \lambda^3 F_1 + \lambda^2\mu F_2 + \lambda\mu^2 F_3 + \mu^3 F_4 = 0, \text{ for } \forall(\lambda : \mu) \in \mathbb{P}^1.$$

So we get the following equations

$$\begin{aligned}
F_1 &:= u_0^3 - (\alpha+1)u_0^2 + \alpha u_0 - u_1^2 + u_3^3 = 0 \\
F_2 &:= 3u_0^2 v_0 - (\alpha+1)2u_0v_0 + \alpha v_0 - 2u_1v_1 + 3u_3^2v_3 - (\beta+1)u_3^2 - u_4^2 = 0 \\
F_3 &:= 3u_0v_0^2 - (\alpha+1)v_0^2 - v_1^2 + 3u_3v_3^2 - (\beta+1)2u_3v_3 + \beta u_3 - 2u_4v_4 = 0 \\
F_4 &:= v_0^3 + v_3^3 - (\beta+1)v_3^2 + \beta v_3 - v_4^2 = 0
\end{aligned}$$

By the implicit function theorem, we can choose the affine coordinates of $F(Y)$ as (u_1, u_4, v_1, v_4).

Let us consider the exceptional divisor of $\nu : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$. Locally, the $\mathbb{Z}_3$ action on $F(Y)$ can be written as

$$\mathbb{Z}_3 \curvearrowright (u_1, u_4, v_1, v_4) \mapsto (u_1, \zeta u_4, \zeta^2 v_1, v_4).$$

The exceptional divisor of $\mathbb{Z}_3$-Hilb(Spec$\mathbb{C}[u_1, u_4, v_1, v_4]/\mathbb{Z}_3$) is given by $E_1 + E_2$, where

$$E_1 = \{(u_1 - \lambda, v_4 - \mu, u_4^2, u_4v_1, v_1^3, au_4 + bv_1^2) | (\lambda, \mu) \in \mathbb{C}^2,\ (a : b) \in \mathbb{P}^1\},$$

$$E_2 = \{(u_1 - \lambda, v_4 - \mu, u_4^2, u_4^3, u_4v_1, v_1^2, cu_4^2 + dv_1) | (\lambda, \mu) \in \mathbb{C}^2,\ (c : d) \in \mathbb{P}^1\}.$$

Here we identify the sub-schemes with their defining ideals. Since the line l corresponds to the origin, when $Z \in E_1$, the ideal sheaf of Z in $G(2,6)$ is

$$I_{(a:b)} = (F_1, F_2, F_3, F_4, u_1, v_4, u_4^2, u_4v_1, v_1^3, au_4 + bv_1^2),\ (a : b) \in \mathbb{P}^1.$$

When $Z \in E_2$, the ideal sheaf of Z in $G(2,6)$ is

$$J_{(c:d)} = (F_1, F_2, F_3, F_4, u_1, v_4, u_4^3, u_4v_1, v_1^3, cu_4^2 + dv_1),\ (c : d) \in \mathbb{P}^1.$$

By the calculation of generators of ideals, we have

$$I_{(a:b)} = (u_0, v_3, v_0, \beta u_3 - v_1^2, u_1, v_4, u_3^2, u_4v_1, v_1^3, au_4 + bv_1^2),\ (a:b) \in \mathbb{P}^1$$

$$J_{(c:d)} = (u_0, u_3, v_3, \alpha v_0 + u_4^2, u_1, v_4, u_4^3, u_4v_1, v_1^2, cu_4^2 + dv_1),\ (c:d) \in \mathbb{P}^1$$

In particular, Z is parametrized by $\mathcal{I} \cup \mathcal{J}$ where $\mathcal{I} = \{I_{(a:b)} | (a:b) \in \mathbb{P}^1\}$ and $\mathcal{J} = \{J_{(c:d)} | (c:d) \in \mathbb{P}^1\}$. Assume that Z is defined by $I_{(a:b)}$. The family of line $\mathcal{L}_Z$ over Z is given by

$$\mathcal{L}_Z = \{(0 : \mu v_1 : \lambda : \lambda u_3 : \lambda u_4 : \mu) | u_3^2 = v_1^3 = \beta u_3 - v_1^2 = au_4 + bv_1^2 = u_4v_1 = 0\}.$$

Since $\beta u_3 - v_1^2 = au_4 + bv_1^2 = 0$, we get an equation $au_4 + b\beta u_3 = 0$. So, the linear space $L_{(a:b)}$ spanned by $\mathcal{L}_Z$ is

$$L_{(a:b)} = \{(x : y : z : u : v : w) \in \mathbb{P}^5 | x = av + b\beta u = 0\}.$$

We remark that $L_{(a:b)}$ is spanned by the tangent line $\{x = 0\} \subset P_C$ of C at p and the line $\{av + b\beta u = 0\} \subset P_D$ passing through $q \in D$.

Assume that Z is defined by $J_{(c:d)}$. In the same way, the linear space $M_{(c:d)}$ spanned by $\mathcal{L}_Z$ is

$$M_{(c:d)} = \{(x : y : z : u : v : w) \in \mathbb{P}^5 | u = dy - c\alpha x = 0\}.$$

$M_{(c:d)}$ is spanned by the line $\{dy - c\alpha x = 0\} \subset P_C$ passing through $p \in C$ and the tangent line $\{u = 0\} \subset P_D$ of D at q. So we can draw the following picture.


$(a : b) = (1 : 0)$
$(a : b)$
$(a : b) = (0 : 1),\ (c : d) = (1 : 0)$
$(c : d)$
$(c : d) = (0 : 1)$
$x = 0$
$C := \{x(x - z)(x - \alpha z) - y^2z = 0\}$
$p = (0 : 0 : 1 : 0 : 0 : 0)$
$D := \{u(u - v)(u - \beta v) - v^2w = 0\}$
$\{(a : b) \in \mathbb{P}^1\}$
$q = (0 : 0 : 0 : 0 : 0 : 1)$
$C := \{x(x - z)(x - \alpha z) - y^2z = 0\}$
$\{(c : d) \in \mathbb{P}^1\}$
$p = (0 : 0 : 1 : 0 : 0 : 0)$
$D := \{u(u - v)(u - \beta v) - v^2w = 0\}$
$u = 0$
$q = (0 : 0 : 0 : 0 : 0 : 1)$

2. By changing coordinates, we may assume that

$$p = (0:1:0),\ f(x,y,z) = x(x-z)(x-\alpha z) - y^2 z,\ \alpha \in \mathbb{C} - \{0,1\}$$

and

$$q = (0:1:0),\ g(u,v,w) = u(u-w)(u-\beta w) - v^2 w,\ \beta \in \mathbb{C} - \{0,1\}.$$

Let U_l be a neighborhood of l in $G(2,6)$. Local coordinates of U_l are given by $(u_0, u_2, u_3, u_5, v_0, v_2, v_3, v_5)$. The corresponding line $\tilde{l}$ to $(u_0, \cdots, v_5)$ is given by

$$(u_0, \cdots, v_5) \leftrightarrow \tilde{l} := \lambda(u_0 : 1 : u_2 : u_3 : 0 : u_5) + \mu(v_0 : 0 : v_2 : v_3 : 1 : v_5)$$

Since $(f+g)(\tilde{l})$ should be zero for each $(\lambda : \mu) \in \mathbb{P}^1$, we have

$$(f+g)(\tilde{l}) = \lambda^3 F_1 + \lambda^2 \mu F_2 + \lambda \mu^2 F_3 + \mu^3 F_4 = 0.$$

Let dF_i be the exterior derivative of F_i $(i = 1, \cdots, 4)$ at the origin. Then we have

$$\begin{pmatrix} dF_1 & dF_2 & dF_3 & dF_4 \end{pmatrix} = \begin{pmatrix} -du_2 & -dv_2 & -du_5 & -dv_5 \end{pmatrix}.$$

The local coordinates around $l \in F(Y)$ is given by (u_0, u_3, v_0, v_3). The $\mathbb{Z}_3$ action on $F(Y)$ is locally given by

$$\mathbb{Z}_3 \curvearrowright F(Y), (u_0, u_3, v_0, v_3) \mapsto (u_0, \zeta u_3, \zeta^2 v_0, v_3).$$

The exceptional divisor of $\mathbb{Z}_3$-Hilb(Spec$\mathbb{C}[u_0, u_3, v_0, v_3]$) is given by $E_1 + E_2$ where

$$E_1 = \{(u_0 - \lambda, v_3 - \mu, u_3^2, u_3 v_0, v_0^3, a u_3 + b v_0^2) | (\lambda, \mu) \in \mathbb{C}^2, (a:b) \in \mathbb{P}^1\},$$

$$E_2 = \{(u_0 - \lambda, v_3 - \mu, u_3^3, u_3 v_0, v_0^2, c u_3^2 + d v_0) | (\lambda, \mu) \in \mathbb{C}^2, (c:d) \in \mathbb{P}^1\}.$$

When $Z \in E_1$, the ideal sheaf of Z in $G(2,6)$ is

$$I_{(a:b)} = (u_2, u_5, v_2, v_5, u_0, v_3, u_3^2, u_3 v_0, v_0^3, a u_3 + b v_0^2),\ (a:b) \in \mathbb{P}^1.$$

When $Z \in E_2$, the ideal sheaf of Z in $G(2,6)$ is

$$J_{(c:d)} = (u_2, u_5, v_2, v_5, u_0, v_3, u_3^3, u_3 v_0, v_0^2, c u_3^2 + d v_0),\ (c:d) \in \mathbb{P}^1.$$

In particular, Z is parametrized by $\mathcal{I} \cup \mathcal{J}$ where $\mathcal{I} = \{I_{(a:b)} | (a:b) \in \mathbb{P}^1\}$ and $\mathcal{J} = \{J_{(c:d)} | (c:d) \in \mathbb{P}^1\}$. Assume that Z is defined by $I_{(a:b)}$. When $(a:b) \neq (1:0)$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = \{(\mu v_0 : \lambda : 0 : \lambda u_3 : \mu : 0) | u_3^2 = u_3 v_0 = v_0^3 = a u_3 + b v_0^2 = 0\}.$$

The linear space $L_{(a:b)}$ spanned by $\mathcal{L}_Z$ is

$$L_{(a:b)} = \{(x:y:z:u:v:w) \in \mathbb{P}^5 | z = w = 0\}.$$

We remark that $L_{(a:b)}$ is spanned by the tangent line $\{z = 0\} \subset P_C$ of C at p and the tangent line $\{w = 0\} \subset P_D$ of D at q. When $(a, b) = (1, 0)$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = \{(\mu v_0 : \lambda : 0 : 0 : \mu : 0) | v_0^3 = 0\}.$$

The linear space $L_{(1:0)}$ spanned by Z is

$$L_{(1:0)} = \{z = u = w = 0\}.$$

The linear space $L_{(1:0)}$ is spanned by the tangent line $\{z = 0\} \subset P_C$ of C at p and the point $q \in D$.

Assume that Z is defined by $J_{(c:d)}$. In the same way, when $(c : d) \neq (0 : 1)$, the linear space $M_{(c:d)}$ spanned by Z is

$$M_{(c:d)} = \{(x:y:z:u:v:w) \in \mathbb{P}^5 | z = w = 0\}.$$

This linear space is spanned by the tangent line $\{z = 0\} \subset P_C$ of C at p and the tangent line $\{w = 0\} \subset P_D$ of D at q. When $(c : d) = (0 : 1)$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = \{(0 : \lambda : 0 : \lambda u_3 : \mu : 0) | u_3^3 = 0\}.$$

So $M_{(0:1)}$ is

$$\{z = w = x = 0\}.$$

The linear space $M_{(0:1)}$ is spanned by the point $p \in C$ and the tangent line $\{w = 0\} \subset P_D$ of D at q.

3. By changing coordinates, we assume that

$$p = (0 : 1 : 0),\ f(x, y, z) = x(x - z)(x - \alpha z) - y^2 z,\ \alpha \in \mathbb{C} - \{0, 1\}$$

and

$$q = (0 : 0 : 1),\ g(u, v, w) = u(u - w)(u - \beta w) - v^2 w,\ \beta \in \mathbb{C} - \{0, 1\}.$$

Let U_l be a neighborhood of l in $G(2, 6)$. Local coordinates of U_l are given by $(u_0, u_2, u_3, u_4, v_0, v_2, v_3, v_4)$. For $(u_0, \cdots, v_4) \in U_l$, the corresponding line $\tilde{l}$ is given by

$$(u_0, \cdots, v_4) \leftrightarrow \tilde{l} := \lambda(u_0 : 1 : u_2 : u_3 : u_4 : 0) + \mu(v_0 : 0 : v_2 : v_3 : v_4 : 1)$$

When $\tilde{l} \in F(Y)$, we have $(f+g)(\tilde{l}) = \lambda^3 F_1 + \lambda^2\mu F_2 + \lambda\mu^2 F_3 + \mu^3 F_4 = 0$, for each $(\lambda, \mu) \in \mathbb{P}^1$. Let dF_i be the exterior derivative of F_i $(i = 1, \cdots, 4)$ at the origin. Then we have

$$\begin{pmatrix} dF_1 & dF_2 & dF_3 & dF_4 \end{pmatrix} = \begin{pmatrix} -du_2 & -dv_2 & \beta du_3 & \beta dv_3 \end{pmatrix}$$

So the local coordinates of $F(Y)$ around l are given by (u_0, u_4, v_0, v_4). The $\mathbb{Z}_3$ action is given by:

$$\mathbb{Z}_3 \curvearrowright F(Y), (u_0, u_4, v_0, v_4) \mapsto (u_0, \zeta u_4, \zeta^2 v_0, v_4).$$

The exceptional divisor of $\mathbb{Z}_3$-Hilb(Spec$\mathbb{C}[u_0, u_4, v_0, v_4]$) is given by $E_1 + E_2$ where

$$E_1 = \{(u_0 - \lambda, v_4 - \mu, u_4^2, u_4v_0, v_0^3, au_4 + bv_0^2) | (\lambda, \mu) \in \mathbb{C}^2, (a:b) \in \mathbb{P}^1\}$$

$$E_2 = \{(u_0 - \lambda, v_4 - \mu, u_4^3, u_4v_0, v_0^2, cu_4^2 + dv_0) | (\lambda, \mu) \in \mathbb{C}^2, (c:d) \in \mathbb{P}^1\}.$$

If $Z \in E_1$ then Z is defined by

$$I_{(a:b)} = (u_2, u_3, v_2, v_3, u_0, v_4, u_4^2, u_4v_0, v_0^3, au_4 + bv_0^2),\ (a:b) \in \mathbb{P}^1$$

If $Z \in E_2$ then the ideal sheaf of Z is

$$J_{(c:d)} = (u_2, u_3, v_3, u_4^2 + v_2, u_0, v_4, u_4^3, u_4v_0, v_0^2, cu_4^2 + dv_0),\ (c:d) \in \mathbb{P}^1.$$

In particular, Z is parametrized by $\mathcal{I} \cup \mathcal{J}$ where $\mathcal{I} = \{I_{(a:b)} | (a:b) \in \mathbb{P}^1\}$ and $\mathcal{J} = \{J_{(c:d)} | (c:d) \in \mathbb{P}^1\}$. Assume that Z is defined by $I_{(a:b)}$. When $(a:b) \neq (1:0)$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = \{(\mu v_0 : \lambda : 0 : 0 : \lambda u_4 : \mu) | u_4^2 = u_4v_0 = v_0^3 = au_4 + bv_0^2 = 0\}.$$

The linear space $L_{(a:b)}$ spanned by $\mathcal{L}_Z$ is

$$L_{(a:b)} = \{z = u = 0\}.$$

This linear space is spanned by the tangent line $\{z = 0\} \subset P_C$ of C at p and the tangent line $\{w = 0\} \subset P_D$ of D at q. When $(a:b) = (1:0)$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = (\mu v_0 : \lambda : 0 : 0 : 0 : \mu) | v_0^3 = 0\}.$$

Then the linear space $L_{(1:0)}$ spanned by $\mathcal{L}_Z$ is

$$L_{(1:0)} = \{z = u = v = 0\}.$$

This linear space is spanned by the tangent line $\{z = 0\} \subset P_C$ of C at p and the point $q \in D$.

Assume that Z is defined by $J_{(c:d)}$. For each $(c : d) \in \mathbb{P}^1$, $\mathcal{L}_Z$ is

$$\mathcal{L}_Z = (\mu v_0 : \lambda : \mu v_2 : 0 : \lambda u_4 : \mu)|u_4^2 + v_2 = u_4^3 = u_4 v_0 = v_0^2 = cu_4^2 + dv_0 = 0\}.$$

Similarly, for each $(c : d) \in \mathbb{P}^1$, the linear space $M_{(c:d)}$ spanned by $\mathcal{L}_Z$ is

$$M_{(c:d)} = \{(x : y : z : u : v : w) \in \mathbb{P}^5 | u = cx - dz = 0\}.$$

The linear space is spanned by the line $\{cx - dz = 0\} \subset P_C$ passing through the point p and the tangent line $\{u = 0\} \subset P_D$ of D at q. $\square$

To describe the indeterminacy of ψ explicitly, we define $Q_{(\mathrm{I})}$ and $Q_{(\mathrm{II})}$ as:

$$Q_{(\mathrm{I})} := \{Z \in \tilde{X}_{\mathbb{Z}_3} | \mathcal{L}_Z \text{ spans } \mathbb{P}^2 \text{ and } \mathcal{L}_Z \ni p \text{ where } p \in C \text{ and } 3p = 0\},$$

and

$$Q_{(\mathrm{II})} := \{Z \in \tilde{X}_{\mathbb{Z}_3} | \mathcal{L}_Z \text{ spans } \mathbb{P}^2 \text{ and } \mathcal{L}_Z \ni q \text{ where } q \in D \text{ and } 3q = 0\}.$$

Let $Q := Q_{(\mathrm{I})} \cup Q_{(\mathrm{II})}$. We shall prove that the indeterminacy of ψ is Q in Theorem 3.5.

Lemma 3.4. *If $\mathcal{L}_Z$ spans $\mathbb{P}^2$, then $\mathcal{L}_Z$ passes through a 3-torsion point of C or D. Furthermore $Q_{(\mathrm{I})}$ is isomorphic to $\{p \in C | 3p = 0\} \times P_D$, and $Q_{(\mathrm{II})}$ is isomorphic to $\{q \in D | 3q = 0\} \times P_C$. In particular, $Q_{(\mathrm{I})}$ and $Q_{(\mathrm{II})}$ are respectively isomorphic to 9 disjoint copies of $\mathbb{P}^2$.*

Proof. We first prove the first assertion. Let l be in $F(Y)$. Since the support of $\mathcal{L}_Z$ is $l \cup \tau(l) \cup \tau^2(l)$, we have to consider the following two cases.

1. $l, \tau(l)$ and $\tau^2(l)$ span $\mathbb{P}^2$.
2. $l = \tau(l) = \tau^2(l)$, that is, l is in the fixed locus of the $\mathbb{Z}_3$ action.

1. Let us consider the first case. Let $P_l := \langle l, \tau(l), \tau^2(l) \rangle$ be the plane spanned by $l, \tau(l)$ and $\tau^2(l)$. Then $\mathbb{Z}_3$ acts on P_l. The eigenvalue of the $\mathbb{Z}_3$ action on $\mathbb{P}^5$ is 1 or $\zeta = \exp(\frac{2\pi\sqrt{-1}}{3})$. So the eigenvalue of the $\mathbb{Z}_3$ action on the plane P_l is also 1 or ζ. Therefore we may assume that, for suitable coordinates of P_l, the group action $\mathbb{Z}_3 \curvearrowright P_l$ is given by

$$\mathbb{Z}_3 \curvearrowright P_l, (x : y : z) \mapsto (x : y : \zeta z).$$

Therefore the fixed locus $\mathrm{Fix}_{\mathbb{Z}_3}(P_l)$ of the $\mathbb{Z}_3$ action on P_l is the disjoint union of a line m_{fix} and a point q_{fix}. Since $\mathrm{Fix}_{\mathbb{Z}_3}(\mathbb{P}^5)$ is $P_C \cup P_D$, we may assume that $m_{\mathrm{fix}} = P_l \cap P_C$, $q_{\mathrm{fix}} = P_l \cap P_D$. Put $p = l \cap m_{\mathrm{fix}}$. Then p is contained in $\mathrm{Fix}_{\mathbb{Z}_3}(Y)$. Since $\mathrm{Fix}_{\mathbb{Z}_3}(Y) = C \cup D$, $p \in C$. We will prove that the point p is a 3-torsion point of C. To prove this, it is enough to show that m_{fix} meets C only at the point p. So we assume that there exists another point p' on $C \cap m_{\mathrm{fix}}$. However $P_l \cap Y$ must be three lines and these three lines meet at exactly one point p. Since p' is in Y, this is contradiction. Therefore p is a 3-torsion point of C. Hence l passes through a 3-torsion point of C.

2. Let us consider the second case. We remark that this case is the limit of the first case. Let Z be in $\tilde{X}_{\mathbb{Z}_3}$ such that the support of $\mathcal{L}_Z$ is l. By the proof of Lemma 3.3, $\mathcal{L}_Z$ which spans $\mathbb{P}^2$ always passes through a 3-torsion point. Hence we get the first assertion.

We shall prove the second assertion for $Q_{(\mathrm{I})}$. Assume that $Z \in Q_{(\mathrm{I})}$. By the above argument, the plane $\langle \mathcal{L}_Z \rangle$ spanned by $\mathcal{L}_Z$ intersect P_D at one point and $\langle \mathcal{L}_Z \rangle \cap P_C$ is the tangent line of C at a 3-torsion point. Conversely we choose a 3-torsion point p of C and a point q from the plane P_D. Let l_C be the tangent line of C at p, and $\langle l_C, q \rangle$ the plane spanned by l_C and q. There uniquely exists $Z \in \tilde{X}_{\mathbb{Z}_3}$ such that $\mathcal{L}_Z$ spans $\langle l_C, q \rangle$. This correspondence gives isomorphism between $Q_{(\mathrm{I})}$ and $\{p \in C | 3p = 0\} \times P_D$. Therefore $Q_{(\mathrm{I})}$ is isomorphic to $\{p \in C | 3p = 0\} \times P_D$. Similarly we can show that $Q_{(\mathrm{II})}$ is isomorphic to $\{q \in D | 3q = 0\} \times P_C$. $\square$

To describe the indeterminacy of $\varphi : K^2(C \times D) \dashrightarrow \tilde{X}_{\mathbb{Z}_3}$, we put

$$\mathbb{P}_{(\mathrm{I})} := \Big\{ \{(p, q_1), (p, q_2), (p, q_3)\} \in K^2(C \times D) \Big| 3p = 0 \Big\},$$

and

$$\mathbb{P}_{(\mathrm{II})} := \Big\{ \{(p_1, q), (p_2, q), (p_3, q)\} \in K^2(C \times D) \Big| 3q = 0 \Big\}.$$

We remark that

$$\mathbb{P}_{(\mathrm{I})} \cong \{p \in C | 3p = 0\} \times P_D^{\vee},$$

and

$$\mathbb{P}_{(\mathrm{II})} \cong \{p \in C | 3p = 0\} \times P_D^{\vee},$$

where $P_D^{\vee}$ (resp. $P_C^{\vee}$) is the dual of P_D (resp. P_C). Moreover, $\mathbb{P}_{(\mathrm{I})}$ and $\mathbb{P}_{(\mathrm{II})}$ are respectively isomorphic to 9 disjoint copies of $\mathbb{P}^2$.

Theorem 3.5. *The birational map $\psi : \tilde{X}_{\mathbb{Z}_3} \dashrightarrow K^2(C \times D)$ is decomposed into the Mukai flop on Q, and φ is decomposed into the Mukai flop on $\mathbb{P}_{(\mathrm{I})} \cup \mathbb{P}_{(\mathrm{II})}$. In particular, the indeterminacy of ψ is Q, and the indeterminacy of φ is $\mathbb{P}_{(\mathrm{I})} \cup \mathbb{P}_{(\mathrm{II})}$.*

The proof of Theorem 3.5 is long. Before starting the proof, we would like to explain the strategy of it.

Step 1. We will blow up $\tilde{X}_{\mathbb{Z}_3}$ along Q. We denote it by $\pi : W \to \tilde{X}_{\mathbb{Z}_3}$. Now let us consider the situation when an irreducible symplectic 4-fold X contains P which is isomorphic to $\mathbb{P}^2$. Since $\mathbb{P}^2$ is a Lagrangian manifold, the normal bundle $N_{P/X}$ of P in X is isomorphic to the cotangent bundle Ω_P^1 of P. Let $B_P X \to X$ be the blow up along P. Then the exceptional locus is isomorphic to

$$B_P X \supset \Gamma := \{(x, H) \in P \times P^{\vee} | x \in H\} \to P \subset X,$$

where $P^{\vee}$ is the dual of P. By Fujiki-Nakano contraction theorem, there exists a birational contraction morphism $B_P X \to X^+$ such that the exceptional locus is isomorphic to $\Gamma \to P^{\vee}$. So, we will blow down W in another direction $\pi^+ : W \to W^+$. Here W^+ is a Moishezon 4-fold. We shall prove that W^+ is actually a projective manifold in Proposition 3.11.

Step 2. We will define a morphism $\mu' : W \to \bar{K}^2(C \times D)$. We will prove that μ' factors through W^+. Namely there is a map $\mu^+ : W^+ \to \bar{K}^2(C \times D)$ such that $\mu^+ \circ \pi^+ = \mu'$.

Step 3. We will prove that W^+ is isomorphic to $K^2(C \times D)$. The following diagram illustrates three steps above.

$$\begin{tikzcd}
& W \arrow[dl, "\pi"'] \arrow[dr, "\pi^+"] \arrow[ddr, "\mu'"'] & & \\
\tilde{X}_{\mathbb{Z}_3} \arrow[rr, dashed] & & W^+ \arrow[d, "\mu^+"] & K^2(C \times D) \arrow[dl, "\mu"'] \\
& & \bar{K}^2(C \times D) &
\end{tikzcd}$$

Let us consider Step 1. We first construct the blow up $\pi : W \to \tilde{X}_{\mathbb{Z}_3}$.

Proposition 3.6. *Let $G(4, 6)$ be the Grassmann manifold which parametrizes 4 dimensional linear subspaces of $\mathbb{C}^6$. We define W by*

$$W := \{(Z, L) \in \tilde{X}_{\mathbb{Z}_3} \times G(4, 6) | \mathcal{L}_Z \subset L, \quad \text{both } L \cap P_C \text{ and } L \cap P_D \text{ are lines}\}.$$

Then the natural projection $\pi : W \to \tilde{X}_{\mathbb{Z}_3}$ is the blow up along Q.

Proof. Let $\pi' : B_Q\tilde{X}_{\mathbb{Z}_3} \to \tilde{X}_{\mathbb{Z}_3}$ be the blow up of $\tilde{X}_{\mathbb{Z}_3}$ along Q. Let $\mathrm{Exc}(\pi')$ be the exceptional divisor of π' and $\mathrm{Exc}(\pi)$ the exceptional divisor of π.

We will define the morphism $f : B_Q\tilde{X}_{\mathbb{Z}_3} \to W$. Outside of the exceptional divisors, we define f by $f := \pi^{-1}\circ\pi'$. On $\mathrm{Exc}(\pi')$, we define f in the following way. On $(\pi')^{-1}(Q_{(\mathrm{I})})$, the morphism π' is defined as follows:

$$\begin{array}{rcl} \{(p,x,H) \in C \times P_D \times P_D^{\vee} | 3p = 0, x \in H\} & \overset{\pi'}{\to} & \{(p,x) \in C \times P_D | 3p = 0\} \\ \pi'(p,x,H) & = & (p,x) \end{array}$$

By Lemma 3.4, (p,x) determine $Z_{(p,x)} \in Q_{(\mathrm{I})}$ uniquely. Here let $l_{C,p}$ be the tangent line of C at p. Hence we define a morphism $f : \mathrm{Exc}(\pi') \to \mathrm{Exc}(\pi)$ by $f(p,x,H) = (Z_{(p,x)}, \langle l_{C,p}, H\rangle)$ where $\langle l_{C,p}, H\rangle$ is the projective linear space spanned by $l_{C,p}$ and H. In the same way, we can define f on $(\pi')^{-1}(Q_{(\mathrm{II})})$. Since f is bijective, W is the blow up along Q. □

As we remarked, by Fujiki-Nakano contraction theorem, there exists a contraction morphism $\pi^+ : W \to W^+$. Note that on $\mathrm{Exc}(\pi^+)$, π^+ can be identified as

$$W \supset \mathrm{Exc}(\pi^+) \ni (Z,L) \to L \in G(4,6).$$

We choose $w \in \pi^+(\mathrm{Exc}(\pi^+))$ and fix it. We remark that for each $(Z,L) \in (\pi^+)^{-1}(w)$, L is uniquely determined by w.

Next let us consider Step 2. We first define $\mu' : W \to \bar{K}^2(C \times D)$ by

$$\mu'(Z,L) := \{(p_i,q_i)\}_{i=1}^3 \text{ where } \langle p_i,q_i\rangle \subset L \cap Y,\ \langle p_i,q_i\rangle \cap \mathcal{L}_Z \neq \emptyset.$$

The map μ' is well-defined.

Lemma 3.7. *The map μ' factors through W^+, that is, there is a map $\mu^+ : W^+ \to \bar{K}^2(C \times D)$ such that $\mu^+ \circ \pi^+ = \mu'$.*

Proof. Let w be a point of W^+. We define μ^+ in the following way:

$$\mu^+(w) := \mu'\big((\pi^+)^{-1}(w)\big).$$

If μ^+ is well-defined then the assertion of Lemma 3.7 is clear. So, we will prove that $\mu'\big((\pi^+)^{-1}(w)\big)$ is independent of the fiber $(\pi^+)^{-1}(w)$. If w is not in $\pi^+(\pi^{-1}(Q))$, then $(\pi^+)^{-1}(w) \cap \pi^{-1}(Q) = \emptyset$. Since the fiber is unique, μ^+ is well-defined.

Assume that w is in $\pi^+(\pi^{-1}(Q))$. Let $(Z,L) \in (\pi^+)^{-1}(w)$. As we remarked, the linear space L is uniquely determined by w. Then exactly one of the following two cases will happen.

1. $L \cap P_C$ is a tangent line of C at a 3-torsion point.

2. $L \cap P_D$ is a tangent line of D at a 3-torsion point.

We may assume that the first case occurs. By Proposition 3.6 $(\pi^+)^{-1}(w)$ is parametrized by the line $L \cap P_D$. Assume that $L \cap P_D \cap D$ consists of three points $\{q_1, q_2, q_3\}$. For each $(Z, L) \in (\pi^+)^{-1}(w)$,

$$\mu'(Z, L) = \{(p, q_1), (p, q_2), (p, q_3)\} \in \bar{K}^2(C \times D).$$

Therefore the map μ^+ is well-defined. □

We consider Step 3 to complete the proof of Theorem 3.5. We will prove that the map μ^+ is isomorphism on the smooth locus of $\bar{K}^2(C \times D)$ in the following Lemma.

Lemma 3.8. *The map μ^+ is an isomorphism on the smooth locus of $\bar{K}^2(C \times D)$.*

Proof. Take $\{(p_i, q_i)\}_{i=1}^3$ from the smooth locus of $\bar{K}^2(C \times D)$. We will prove that the fiber $(\mu^+)^{-1}(\{(p_i, q_i)\})$ (for short $\{(p_i, q_i)\}_{\mu^+}$) is one point. Let $l_C \subset P_C$ (resp. $l_D \subset P_D$) be the line passing through three points $\{p_i\}_{i=1}^3$ (resp. $\{q_i\}_{i=1}^3$). We have to consider the fibers of μ^+ in the following cases.

1. $\bar{K}^2(C \times D) \ni \{(p_i, q_i)\}_{i=1}^3$ where $p_i \neq p_j$ $(i \neq j)$, and $q_i \neq q_j$ $(i \neq j)$.
2. $\bar{K}^2(C \times D) \ni \{(p_i, q_i)\}_{i=1}^3$ where $p_1 = p_2$, $p_1 \neq p_3$ and $q_i \neq q_j$ $(i \neq j)$.
3. $\bar{K}^2(C \times D) \ni \{(p_i, q_i)\}_{i=1}^3$ where $p_1 = p_2$, $p_1 \neq p_3$ and $q_2 = q_3$, $q_2 \neq q_1$.
4. $\bar{K}^2(C \times D) \ni \{(p_i, q_i)\}_{i=1}^3$ where $p_1 = p_2 = p_3 =: p$, and $q_i \neq q_j$ $(i \neq j)$.

1. Let us consider the first case. In this case, the fiber $\{(p_i, q_i)\}_{\mu^+}$ is isomorphic to the fiber $(\mu')^{-1}(\{(p_i, q_i)\})$ (for short $\{(p_i, q_i)\}_{\mu'}$). So assume that $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$. The linear space L is spanned by l_C and L_D, and the cubic surface $L \cap Y$ is smooth. As we wrote in Proposition 3.2, Z consists of three lines l, $\tau(l)$ and $\tau^2(l)$. In particular Z is uniquely determined. Hence the fiber is one point.

2 and 3. Notice that the case 3 is the limit of the case 2. Since $\{(p_i, q_i)\}_{\mu'} \cap \pi^{-1}(Q) = \emptyset$, the fiber $\{(p_i, q_i)\}_{\mu'}$ is isomorphic to the fiber $\{(p_i, q_i)\}_{\mu^+}$. Let $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$. The linear space L is spanned by l_C and l_D. Let us consider the cubic surface $L \cap Y$. There is a line $l \subset L \cap Y$ which intersect three lines $\{\langle p_i, q_i \rangle\}_{i=1}^3$. In both case 2 and 3, the line l passes through p_1 and q_3. Notice that the line l is in the fixed locus of the $\mathbb{Z}_3$ action on $F(Y)$. Since there uniquely exists Z such that $\mathcal{L}_Z$ spans L, the fiber is one point.

4. Let us consider the fourth case. Assume that $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$. By the similar argument in the proof of Lemma 3.7, $\{(p_i, q_i)\}_{\mu'}$ is parametrized by the line l_D. Since $\{(p_i, q_i)\}_{\mu'}$ is contracted to a point by π^+, the fiber $\{(p_i, q_i)\}_{\mu^+}$ is one point. □

We will use the following result to complete Step 3.

Proposition 3.9 ([F-N]). *Let Y be a locally $\mathbb{Q}$ factorial symplectic variety, and suppose that there is a projective crepant resolution of Y: $\pi : X \to Y$. If the exceptional divisor of π is irreducible, then π is a unique crepant resolution, that is, if there is another crepant resolution $\pi' : X' \to Y$ then the natural birational map $\pi' \circ \pi^{-1}$ is an isomorphism.*

Since the exceptional divisor of $\mu : K^2(C\times D) \to \bar{K}^2(C\times D)$ is irreducible, it is enough to show that W^+ is projective. We will observe the fiber of singular points of $\bar{K}^2(C \times D)$ by μ^+ and prove that W^+ is projective

Proposition 3.10. *Let E be the exceptional divisor of $\mu^+ : W^+ \to \bar{K}^2(C \times D)$. Then $-E$ is μ^+-ample. In particular W^+ is a projective manifold.*

Proof. We first consider the fiber of singular points by μ^+. By Lemma 3.8, the exceptional divisor of π^+ is contained in the inverse image of the singular locus of $\bar{K}^2(C \times D)$. We will use the same notation in the proof of Lemma 3.8. By symmetry we have to consider the following three cases.

1. $\{(p_i, q_i)\}_{i=1}^3 \in \bar{K}^2(C \times D)$. $p_1 = p_2 =: p$, $p_1 \neq p_3 = -2p$ and $q_1 = q_2 =: q$, $q_1 \neq q_3 = -2q$.
2. $\{(p_i, q_i)\}_{i=1}^3 \in \bar{K}^2(C \times D)$. $p_1 = p_2 = p_3 =: p$ and $q_1 = q_2 =: q$, $q_1 \neq q_3 = -2q$.
3. $\{(p_i, q_i)\}_{i=1}^3 \in \bar{K}^2(C \times D)$. $p_1 = p_2 = p_3 =: p$ and $q_1 = q_2 = q_3 =: q$

1. Let us consider the first case. We claim that the fiber $\{(p_i, q_i)\}_{\mu^+}$ is isomorphic to $\mathbb{P}^1$. Since $\{(p_i, q_i)\}_{\mu'} \cap \pi^{-1}(Q) = \emptyset$, we have $\{(p_i, q_i)\}_{\mu^+} \cong \{(p_i, q_i)\}_{\mu'}$. So, let $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$. Since the linear space L is spanned by the line l_C and l_D, L is unique. There are infinitely many lines in the cubic surface $L \cap Y$ which intersect with two lines $\langle p, q\rangle$ and $\langle -2p, -2q\rangle$. Here we can draw the following picture of the configuration of lines in $L \cap Y$.



The set of lines can be parametrized by the line $\langle p, q\rangle$, that is, the line l in $L \cap Y$ is uniquely determined by $t \in \langle p, q\rangle$. The cubic surface has singular points along the line $\langle p, q\rangle$.

Notice that $\mathbb{Z}_3$ acts on the line $\langle p, q\rangle$. The fiber $\{(p_i, q_i)\}_{\mu'}$ is isomorphic to $\langle p, q\rangle/\mathbb{Z}_3$. Indeed let $l_t \ \subset L \cap Y$ be the line passing through $t \in \langle p, q\rangle$. When $t \in \langle p, q\rangle - \{p, q\}$, l_t is not in the fixed locus of the $\mathbb{Z}_3$ action on $F(Y)$. Hence there exists uniquely Z in $\tilde{X}_{\mathbb{Z}_3}$ such that the support of $\mathcal{L}_Z$ is $l_t \cup \tau(l_t) \cup \tau^2(l_t)$. When $t = p$, l_p passes through p and $-2q$. Notice that l_p is in the fixed locus of the $\mathbb{Z}_3$ action on $F(Y)$. By Lemma 3.3, there uniquely exists $Z \in \tilde{X}_{\mathbb{Z}_3}$ such that the support of $\mathcal{L}_Z$ is l_p and $\mathcal{L}_Z$ spans L. Similarly, if $t = q$, then the line l_q passes through $-2p$ and q. By L, Z can be determined uniquely. So $\{(p_i, q_i)\}_{\mu^+}$ is isomorphic to $\langle p, q\rangle/\mathbb{Z}_3$.

2. Let us consider the second case. Recall the notation $\mathcal{I}$ defined in third case of the proof for Lemma 3.3. We will prove that the fiber $\{(p_i, q_i)\}_{\mu^+}$ is parametrized by $\mathcal{I}$. Let $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$. Then L is spanned by l_C and l_D. Furthermore, the cubic surface $L \cap Y$ is given by $\{(x : y : z : w) \in L | y^3 + z^2 w = 0\}$ for suitable homogeneous coordinates of L. The point $(1 : 0 : 0 : 0) \in L$ correspond to p. Notice that all lines contained in the cubic surface $L \cap Y$ pass through p.

We claim that the fiber $\{(p_i, q_i)\}_{\mu'}$ is isomorphic to the tree $\mathcal{I} \cup l_D$ of two projective line $\mathcal{I}$ and l_D. Let P_t be the plane spanned by l_C and $t \in l_D$. If $t \in l_D - \{q, -2q\}$, then $P_t \cap Y$ is not in the fixed locus. So we can uniquely determine Z such that the support of $\mathcal{L}_Z$ is $P_t \cap Y$. When $t = -2q$, $P_t \cap Y$ is the line $\langle p, -2q\rangle$ passing through p and $-2q$. There uniquely exists Z such that the support of $\mathcal{L}_Z$ is the line $\langle p, -2q\rangle$ and $\mathcal{L}_Z$ spans L. When $t = q$, $P_t \cap Y$ is the line $\langle p, q\rangle$. Here recall that $I_{(a:b)} \in \mathcal{I}$ is the ideal defined in the third case in the proof of Lemma 3.3. Let $Z_{(a:b)} \in \tilde{X}_{\mathbb{Z}_3}$ be the point defined by $I_{(a:b)}$. For each $I_{(a:b)} \in \mathcal{I}$, $(Z_{(a:b)}, L)$ is in the fiber $\{(p_i, q_i)\}_{\mu'}$. So in the second case, we can not uniquely determine Z. Therefore, $\{(p_i, q_i)\}_{\mu'}$ is isomorphic to $\mathcal{I} \cup l_D$.

Since the line l_D is contracted to a point by $\pi^+ : W \to W^+$, the fiber $\{(p_i, q_i)\}_{\mu^+}$ is isomorphic to $\mathcal{I}$. We draw the picture of the "blow down".



3. Let us consider the third case. When $(Z, L) \in \{(p_i, q_i)\}_{\mu'}$, the linear space L is spanned by l_C and l_D. Moreover the cubic surface $L \cap Y$ is three planes P_1, P_2 and P_3. Each plane is sent to another plane by the $\mathbb{Z}_3$ action. Three planes share the line $\langle p, q\rangle$. So we choose a plane P_1 and fix it.



For each $t \in \langle p, q\rangle$, the set of lines passing through t in P_1 is parametrized by $\mathbb{P}^1$. This parameter means direction of the line passing through t. So we call this parameter $D_t(s)$ where $s \in \mathbb{P}^1$. For each t, there is a special direction $D_t(s_0)$ which represent the line $\langle p, q\rangle$. Notice that $\langle p, q\rangle$ is in the fixed locus of the $\mathbb{Z}_3$ action on $F(Y)$. Here recall that $\mathcal{I}$ and $\mathcal{J}$ defined in the second case of the poof for Lemma 3.3 and let $I_{(a:b)} \in \mathcal{I}$ and $J_{(c:d)} \in \mathcal{J}$. If we consider the blow up $\tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$, then the tree $\mathcal{I} \cup \mathcal{J}$ appears as the exceptional curves. Now let us consider the limit $\lim_{s\to s_0} D_t(s)$ for $t \in \langle p, q\rangle - \{p, q\}$. This limit is independent of t and $\lim_{s\to s_0} D_t(s)$ coincides with the intersection $I \cap \mathcal{J}$. If we flop $\tilde{X}_{\mathbb{Z}_3} \dashrightarrow W^+$, then $D_p(s)$ and $D_q(s)$ are contracted to points. So the fiber $\{(p_i, q_i)\}_{\mu^+}$ is the contraction of Hirtzebruch surface $F_n \to \tilde{F}_n$ by $(-n)$-curve. In particular, the Picard number of the surface $\{(p_i, q_i)\}_{\mu^+}$ is 1. We can explain these by the following picture.



Let E be the exceptional divisor of μ^+. E has a fibration over the singular points of $\bar{K}^2(C \times D)$. A general fiber of the fibration is $\mathbb{P}^1$ and there are 81 singular fibers $\tilde{F}_n$. Since the Picard number of $\tilde{F}_n$ is 1, all curves contracted by μ^+ are algebraically equivalent to a general fiber

$$\xi = (\mu^+)^{-1}(\{(p,q),(p,q),(-2p,-2q)\}).$$

Since the normal bundle of ξ is trivial, the intersection number (ξ, E) is -2. Therefore $-E$ is μ^+ ample. $\square$

3.2 Birational symplectic models of $\tilde{X}_{\mathbb{Z}_3}$

Finally let us consider other birational symplectic models of $\tilde{X}_{\mathbb{Z}_3}$ which are different from $K^2(C \times D)$. We define divisors needed later. Let E_1 be the irreducible component of $\mathrm{Exc}(\nu) : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$ such that E_1 meets $Q_{(\mathrm{I})}$, and E_1 does not meet $Q_{(\mathrm{II})}$. Similarly, let E_2 be the irreducible component of $\mathrm{Exc}(\nu)$ such that E_2 meets $Q_{(\mathrm{II})}$, and E_2 does not meet $Q_{(\mathrm{I})}$. We remark that $\mathrm{Exc}(\nu)$ is $E_1 + E_2$. Here let $K^2(C \times D) \dashrightarrow X_{\mathrm{I}}$ be the Mukai flop on $\mathbb{P}_{(\mathrm{I})}$, and $K^2(C \times D) \dashrightarrow X_{\mathrm{II}}$ the Mukai flop on $\mathbb{P}_{(\mathrm{II})}$.

Theorem 3.11. *Assume that C and D are "not" isogenus. Then there are exactly 4 birational models of $\tilde{X}_{\mathbb{Z}_3}$, that is, there are 4 projective irreducible symplectic manifolds $K^2(C \times D)$, $\tilde{X}_{\mathbb{Z}_3}$, X_{I} and X_{II} which are birational to $K^2(C \times D)$. The movable cone of $\tilde{X}_{\mathbb{Z}_3}$ is decomposed into the ample cones of these 4 models in the following way.*



Each codimension 1 face corresponds to the following contraction morphism.

Face	*Morphism*
$\overline{oc}$	*Contraction of* $\mathbb{P}_{(\mathrm{I})}$
$\overline{od}$	*Contraction of* $\mathbb{P}_{(\mathrm{II})}$
$\overline{cd}$	$K^2(C \times D) \to \bar{K}^2(C \times D)$
$\overline{ab}$	*Contraction of* E_1
$\overline{ae}$	*Contraction of* E_2
$\overline{ob}$	*Contraction of* $Q_{(\mathrm{II})}$
$\overline{oe}$	*Contraction of* $Q_{(\mathrm{I})}$
$\overline{bc}$	*Contraction of* $\tilde{E}_1$
$\overline{de}$	*Contraction of* $\tilde{E}_2$

Vertex	*Morphism*
c	$K^2(C \times D) \to K^2(C)$
d	$K^2(C \times D) \to K^2(D)$
o	*Contraction of 18 copies of* $\mathbb{P}^2$
a	$\tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$
b	*contraction of* E_1 *and* $Q_{(\mathrm{II})}$.
e	*contraction of* E_2 *and* $Q_{(\mathrm{I})}$.

Here $\tilde{E}_1$ *is the proper transform of* E_1 *in* X_{I}, $\tilde{E}_2$ *is the proper transform of* E_2 *in* X_{II}.

We will provide the proof into three parts. We always assume that C and D are not isogenus.

Lemma 3.12. *The ample cone of* $K^2(C \times D)$ *is given by the following picture, where* H_1 *(resp.* H_2, $6H_1 + 6H_2 - E$*) is the supporting divisor of the vertex* c *(resp.* d, o*).*



The divisors H_1, H_2 *and* E *will be defined in the proof.*

Proof. Since C and D are not isogenus, the Picard number of $K^2(C \times D)$ is 3. Let E be the exceptional divisor of $\mu : K^2(C \times D) \to \bar{K}^2(C \times D)$. We have the following commutative diagram.

$$\begin{tikzcd}
 & K^2(C \times D) \arrow[dl, "\pi_1"'] \arrow[d, "\mu"] \arrow[dr, "\pi_2"] & \\
K^2(C) & \bar{K}^2(C \times D) \arrow[l, "\bar{\pi}_1"'] \arrow[r, "\bar{\pi}_2"] & K^2(D)
\end{tikzcd}$$

Let H_C (resp. H_D) be the pull back of the tautological line bundle $\mathcal{O}_{\mathbb{P}^2}(1)$ by the isomorphism $K^2(C) \to \mathbb{P}^2$ (resp. $K^2(D) \to \mathbb{P}^2$). We define H_1 and H_2 by:

$$H_1 := \pi_1^*(H_C),\ H_2 := \pi_2^*(H_D).$$

Let x be a line in $\mathbb{P}_{(\mathrm{I})}$, y a line in $\mathbb{P}_{(\mathrm{II})}$ and z a rational curve which can be contracted by μ. Write $\mu(z) = \{(p,q),(p,q),(-2p,-2q)\}$ and assume that neither p nor q are 3-torsion points.

Since x is contracted by π_1, the intersection number (x, H_1) is zero. Since π_2 is an isomorphism on one connected component of $\mathbb{P}_{(\mathrm{I})}$, we have $(x, H_2) = 1$. In the same way, $(y, H_2) = 0$, and $(y, H_1) = 1$. Since z is contracted by π_1 and π_2, we have $(z, H_1) = (z, H_2) = 0$. Since the normal bundle of z in E is trivial and z is isomorphic to $\mathbb{P}^1$, $(z, E) = -2$. The intersection $E \cap \mathbb{P}_{(\mathrm{I})}$ is given by

$$E \cap \mathbb{P}_{(\mathrm{I})} \quad = \quad \big\{\{(p,q), (p,q), (p,-2q)\} \in K^2(C \times D) \big| 3p = 0, q \in D\big\}.$$

So, $E \cap \mathbb{P}_{(\mathrm{I})}$ is isomorphic to 9 copies of the curve

$$\big\{\{q, q, -2q\} \in K^2(D) | q \in D\big\}.$$

This curve is isomorphic to the dual curve $D^\vee$ of D. Since $\deg D^\vee = 6$, $(x, E) = 6$. Similarly, we have $(y, E) = 6$. From these arguments, we have the the following table of intersection numbers.

	H_1	H_2	E
x	0	1	6
y	1	0	6
z	0	0	-2

Clearly both H_1 and H_2 are nef. We will prove that $6H_1 + 6H_1 - E$ is nef. Notice that $6H_1$ (resp. $6H_2$) is lineally equivalent to $\pi_1^* C^\vee$ (resp. $\pi_2^* D^\vee$). Since $\bar{\pi}_1^* C^\vee$ contains the singular locus of $\bar{K}^2(C \times D)$, $\pi_1^* C^\vee$ has two irreducible components. Let F_1 be the proper transform of $\bar{\pi}_1^* C^\vee$ and F_2 the proper transform of $\bar{\pi}_2^* D^\vee$. Assume that $\pi_1^* C^\vee \sim aF_1 + bE$ for some numbers a and b. Since $(z, F_1) = 1$ and $(y, F_1) = 0$, we have $\pi_1^* C^\vee = 2F_1 + E$. Similarly, $\pi_2^* D^\vee \sim 2F_2 + E$. Assume that $6H_1 + 6H_2 - E$ is not nef. Then there is a curve ξ such that $(\xi, 6H_1 + 6H_2 - E) < 0$. Since $6H_1 + 6H_2 - E$ is linearly equivalent to the following divisors, ξ should be contained in $F_1 \cap F_2$.

$$\begin{aligned} 6H_1 + 6H_2 - E &\sim \pi_1^* C^\vee + 6H_2 - E = 2F_1 + 6H_2 \\ &\sim 6H_1 + \pi_2^* D^\vee - E = 6H_1 + 2F_2. \end{aligned}$$

By using the cone theorem, we can contract ξ. Theorem 1 in [Kaw] says that the exceptional locus is covered by rational curves. Since $F_1 \cap F_2$ is isomorphic to $C^\vee \times D^\vee$, $F_1 \cap F_2$ does not contain the rational curve. So, $6H_1 + 6H_2 - E$ is nef.

By the table of intersection numbers, the face $\overline{oc}$ (resp. $\overline{od}$ and $\overline{cd}$) defines the contraction of x (resp. y and z). □

Lemma 3.13. *The ample cone of* $\tilde{X}_{\mathbb{Z}_3}$ *is given by:*



.

Each supporting divisor will be defined in the proof.

Proof. Let $K^2(C\times D)\to X_0$ be the contraction of $\mathbb{P}_{(\mathrm{I})}\cup\mathbb{P}_{(\mathrm{II})}$. By Theorem 3.5, there exists a projective morphism $\phi:\tilde{X}_{\mathbb{Z}_3}\to X_0$, and we have the following diagram:

$$\begin{tikzcd}
 & W \arrow[dl, "\pi^+"'] \arrow[dr, "\pi"] & \\
K^2(C\times D) \arrow[rr, dashed, "\text{Mukaiflop}", "\varphi"'] \arrow[dr] & & \tilde{X}_{\mathbb{Z}_3} \arrow[dl, "\phi"] \\
 & X_0 &
\end{tikzcd}$$

Let H_{X_0} be an ample divisor of X_0. Then we may assume that $\phi^*H_{X_0}$ is numerically equivalent to the proper transform of $6H_1+6H_2-E$. Let x be a rational curve in E_1 which is contracted by ν, y a rational curve in E_2 which is contracted by ν, z a line in $Q_{(\mathrm{I})}$, and w a line in $Q_{(\mathrm{II})}$.

Clearly $(x,E_2)=(y,E_1)=1$. Since the normal bundle of x in E_1 (resp. y in E_2) is trivial, $(x,E_1)=-2$ (resp. $(y,E_2)=-2$). By the definition of X_0, we have $(z,\phi^*H_{X_0})=(w,\phi^*H_{X_0})=0$. Since $Q_{(\mathrm{I})}\cap E_1$ is isomorphic to D, we have $(z,E_1)=3$. Since $Q_{(\mathrm{I})}\cap E_2=\emptyset$, we have $(z,E_2)=0$. Similarly, we have $(w,E_2)=3$ and $(w,E_1)=0$. Recall divisors F_1, F_2 and E of $K^2(C\times D)$ defined in the proof for Lemma 3.12. Let $\tilde{F_1}$, $\tilde{F_2}$ and $\tilde{E}$ be a proper transform of F_1, F_2 and E respectively. By Lemma 3.3, we have $\tilde{F_1}=E_1$ and $\tilde{F_2}=E_2$. Here, we will prove that $(x,\phi^*H_{X_0})=2$. Let $Z\in\tilde{X}_{\mathbb{Z}_3}$ be in the curve x. For each Z, the support of $\mathcal{L}_Z$ is a line $\langle p_0,q_0\rangle$ passing through $p_0\in C$ and $q_0\in D$. Then $x\cap\tilde{E}$ is identified with 4 points $\{q\in D|-2q=q_0\}$. Hence $(x,\tilde{E})=4$. Since $6H_1+6H_2-E\sim 2F_1+2F_2+E$, we have the following equation:

$$\begin{aligned}4=(x,\tilde{E}) &= (x,\phi^*H_{X_0}-2\tilde{F_1}-2\tilde{F_2})\\ &= (x,\phi^*H_{X_0})-2(x,E_1)-2(x,E_2).\end{aligned}$$

Hence we have $(x,\phi^*H_{X_0})=2$. In the same way, we have $(y,\phi^*H_{X_0})=2$. Therefore we get the following table of intersection numbers.

	$\phi^*H_{X_0}$	E_1	E_2
x	2	-2	1
y	2	1	-2
z	0	3	0
w	0	0	3

We will prove that the divisor $\phi^*H_{X_0} + 2E_1 + 2E_2$ defines the morphism $\nu : \tilde{X}_{\mathbb{Z}_3} \to X_{\mathbb{Z}_3}$. Suppose that the vertex a is generated by ν^*H, where H is an ample divisor of $X_{\mathbb{Z}_3}$. Since the Picard number of $\tilde{X}_{\mathbb{Z}_3}$ is 3, we put $\nu^*H = a\phi^*H_{X_0} + bE_1 + cE_2$ for some numbers (a, b, c). Since both $(x, \nu^*H) = (y, \nu^*H) = 0$, we have $b = c = 2a$.

We will prove that both $\phi^*H_{X_0} + E_1$ and $\phi^*H_{X_0} + E_2$ are nef. From symmetry it is enough to prove it only for $\phi^*H_{X_0} + E_1$. Since $\phi^*H_{X_0}$ is nef, the curve ξ such that $(\xi, \phi^*H_{X_0} + E_1) < 0$ should be contained in E_1. We can contract ξ by the cone theorem, and Theorem 1 in [Kaw] says that such a curve must be rational. Since E_1 is $\mathbb{P}^1$ bundle over $C \times D$, it is enough to check that $(x, \phi^*H_{X_0} + E_1) \geq 0$. By the above table of intersection number, we have

$$(x, \phi^*H_{X_0} + E_1) = 0.$$

Therefore $\phi^*H_{X_0} + E_1$ is nef.

By the table of intersection numbers, we have the correspondence between vertices and contraction morphisms in Theorem 3.11. □

To complete the proof of Theorem 3.11, we consider the ample cone of X_{I} and X_{II}. By symmetry, it is enough to prove it for X_{I}.

Lemma 3.14. *The ample cones of* X_{I} *and* X_{II} *are given by*:



Proof. Throughout the proof,we use the same notations used in the proof of Lemma 3.13. We only prove that the ample cone of $X_{(\mathrm{I})}$ is a "tetrahedron". In $\tilde{X}_{\mathbb{Z}_3}$, the vertex b corresponds to the contraction of $E_1 \cup Q_{(\mathrm{II})}$. We denote by $\tilde{E}_1 \subset X_{\mathrm{I}}$ the proper transform of E_1 and by $\tilde{x} \subset X_{\mathrm{I}}$ the proper transform of x. Let $K^2(C \times D) \to Z_{\mathrm{I}}$ be a contraction of $\mathbb{P}_{(\mathrm{I})}$. By the construction of

X_{I}, there exists a birational contraction morphism $\psi_{\mathrm{I}} : X_{\mathrm{I}} \to K^2(C)$ such that the following diagram commutes.

$$\begin{tikzcd}
X_{\mathrm{I}} \arrow[dr] \arrow[ddr, "\psi_{\mathrm{I}}"'] & & K^2(C \times D) \arrow[ll, dashed, "\text{Mukai flop on } \mathbb{P}_{(\mathrm{I})}"'] \arrow[dl] \arrow[ddl, "\pi_1"] \\
& Z_{\mathrm{I}} \arrow[d] & \\
& K^2(C) &
\end{tikzcd}$$

In X_{I}, the vertex c corresponds to the contraction morphism ψ_{I}. Hence it is enough to prove that $\tilde{x}$ is contracted by ψ_{I}. Assume that $Z \in \tilde{X}_{\mathbb{Z}_3}$ is in the curve x. For each $Z \in x$, the support of $\mathcal{L}_Z$ is the line $\langle p, q \rangle$ passing through $p \in C$ ($3p \neq 0$) and $q \in D$ ($3q \neq 0$). Let $\xi \subset K^2(C \times D)$ be the proper transform of $\tilde{x}$. Then the curve $\xi \subset K^2(C \times D)$ is given by:

$$\xi = \big\{ \{(p, q_1), (p, q_2), (-2p, q)\} \in K^2(C \times D) | q_1 + q_2 + q = 0 \big\}.$$

So $\tilde{x}$ is contracted by ψ_{I}. $\square$

4 Second example

Let M be the finite subset of $PGL(5)$ defined by

$$M := \Bigg\{ A := \begin{pmatrix} a_0 & & \\ & \ddots & \\ & & a_5 \end{pmatrix} \in PGL(5) \Bigg| A \text{ is a diagonal matrix,}$$
$$\#\{i | a_i = 1\} = 3, \#\{j | a_j = \zeta\} = 3 \ \cdots \ (*) \Bigg\}$$

Then $\#M = 10$. Let G be the finite group generated by the set M. We define $\tau_i (i = 1, \cdots 4) \in G$ respectively by:

$$\begin{aligned}
\tau_1 &:= \{a_0 = a_1 = a_2 = 1, a_3 = a_4 = a_5 = \zeta\} \\
\tau_2 &:= \{a_0 = a_1 = a_5 = 1, a_2 = a_3 = a_4 = \zeta\} \\
\tau_3 &:= \{a_0 = a_1 = a_2 = a_3 = 1, a_4 = \zeta, a_5 = \zeta^2\} \\
\tau_4 &:= \{a_0 = a_3 = a_4 = a_5 = 1, a_1 = \zeta, a_2 = \zeta^2\}.
\end{aligned}$$

Then τ_1, τ_2, τ_3 and τ_4 are generators of G. So G is isomorphic to $\mathbb{Z}_3^{\oplus 4}$. Let Y be a smooth cubic 4-fold of Fermat type:

$$Y := \{(z_0 : \cdots : z_5) \in \mathbb{P}^5 | z_0^3 + \cdots + z_5^3 = 0\}.$$

Let C be the Fermat type elliptic curve:$\{(x:y:z) \in \mathbb{P}^2 | x^3+y^3+z^3=0\}$. We will consider the induced G-action on $F(Y)$.

Lemma 4.1. *The G-action preserves the symplectic form $\sigma_{F(Y)}$ and $X_G = F(Y)/G$ has a crepant resolution $\tilde{X_G}$.*

Proof. By Lemma 2.3, G preserves the symplectic form $\sigma_{F(Y)}$. We put

$$\alpha_i = \mathrm{diag}(a_0^{(i)}, \cdots, a_5^{(i)}) \in M \ (i = 1, \cdots, 10).$$

We may assume that α_1 is τ_1 and α_2 is τ_2. Each α_i generates the cyclic group of order 3. Let $\mathrm{Fix}(\alpha_{\mathrm{i}})(i=1,\cdots,10)$ be the fixed locus of the action $\langle \alpha_i \rangle \curvearrowright F(Y)$. Then $\mathrm{Fix}(\alpha_i)$ is isomorphic to $C \times C$


$\mathrm{Fix}(\alpha_1) \cong C \times C$
9 copies of 9 points
A_2sing × A_2sing
……
$\mathrm{Fix}(\alpha_1) \cap \mathrm{Fix}(\alpha_2)$
$\mathrm{Fix}(\alpha_1) \cap \mathrm{Fix}(\alpha_{10})$

Each intersection $\mathrm{Fix}(\alpha_i) \cap \mathrm{Fix}(\alpha_j)(i \neq j)$ consists of 9 points. Since the G-action preserves the symplectic form, singularities of X_G is the product of two A_2 singularities at each point of $\mathrm{Fix}(\alpha_i) \cap \mathrm{Fix}(\alpha_j)$. So X_G has a crepant resolution $\tilde{X_G}$. ☐

Since C has a complex multiplication, $\mathbb{Z}_3$ acts on C in the following way:

$$\mathbb{Z}_3 \curvearrowright C, \ x \mapsto \zeta x.$$

We consider the following $\mathbb{Z}_3$ action on $C \times C$:

$$\mathbb{Z}_3 \curvearrowright C \times C, \ (x, y) \mapsto (\zeta x, \zeta^2 y).$$

Let S be the minimal resolution of $C \times C/\mathbb{Z}_3$. Notice that S is a K3 surface.

Theorem 4.2. *Notations being as above, $\tilde{X_G}$ is birational to* $\mathrm{Hilb}^2(S)$

Proof. To distinguish two elliptic curves $\{(z_0 : z_1 : z_2) \in \mathbb{P}^2 | z_0^3 + z_1^3 + z_2^3 = 0\}$ and $\{(z_3 : z_4 : z_5) \in \mathbb{P}^2 | z_3^3 + z_4^3 + z_5^3 = 0\}$, we call them C and D respectively. Let $\mathfrak{S}_3$ be the symmetric group with degree 3. By Theorem 3.5 we have the following birational map:

$$\begin{aligned}\tilde{X}_G \overset{\text{birat.}}{\sim} F(Y)/\langle \tau_1, \tau_2, \tau_3, \tau_4 \rangle &\overset{\text{birat.}}{\sim} \bar{K}^2(C \times D)/\langle \tau_2, \tau_3, \tau_4 \rangle \\ &= \Big((C \times C \times D \times D)/\mathfrak{S}_3\Big)\Big/\langle \tau_2, \tau_3, \tau_4 \rangle.\end{aligned}$$

We will show that $\Big((C \times D)^2/\mathfrak{S}_3\Big)\Big/\langle \tau_2, \tau_3, \tau_4 \rangle$ is birational to $\mathrm{Hilb}^2(S)$. Let x_0(resp. y_0) be a 3-torsion point of C(resp. D) such that $\zeta x_0 = x_0$. The induced actions of τ_2, τ_3 and τ_4 on $(C \times D)^2$ are give by

$$\begin{aligned}\tau_2 \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (\zeta x_1, \zeta x_2, \zeta^2 y_1, \zeta^2 y_2) \\ \tau_3 \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (x_1, x_2, y_1 + y_0, y_2 + y_0) \\ \tau_4 \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (x_1 + x_0, x_2 + x_0, y_1, y_2).\end{aligned}$$

Here $\mathfrak{S}_3$ is generated by

$$\mathfrak{S}_3 = \langle \sigma, \tau | \sigma^3 = 1, \tau^2 = 1, \sigma\tau = \tau\sigma^2 \rangle,$$

and the action $\mathfrak{S}_3$ on $C \times C \times D \times D$ is given by

$$\begin{aligned}\sigma \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (x_2, -x_1 - x_2, y_2, -y_1 - y_2) \\ \tau \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (x_2, x_1, y_2, y_1).\end{aligned}$$

Let us diagonalize the actions $\sigma \curvearrowright C \times C$. We have the following diagram:

$$\begin{tikzcd} C \times C \arrow[r, "f"] \arrow[d, "\sigma"'] & C \times C \arrow[d, "\tilde{\sigma}"] \\ C \times C \arrow[r, "f"] & C \times C, \end{tikzcd}$$

where the matrix representations of σ, f and $\tilde{\sigma}$ are respectively

$$\sigma = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}, \ f = \begin{pmatrix} \zeta^2 & -1 \\ -\zeta & 1 \end{pmatrix}, \ \tilde{\sigma} = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^2 \end{pmatrix}.$$

For τ, we have the following diagram:

$$\begin{tikzcd} C \times C \arrow[r, "f"] \arrow[d, "\tau"'] & C \times C \arrow[d, "\tilde{\tau}"] \\ C \times C \arrow[r, "f"] & C \times C, \end{tikzcd}$$

where the matrix representations of τ and $\tilde{\tau}$ are respectively

$$\tau = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \tilde{\tau} = \begin{pmatrix} 0 & \zeta \\ \zeta^2 & 0 \end{pmatrix}.$$

We remark that $(x_1, x_2) \in C \times C$ is in the kernel of f if and only if $\zeta^2 x_1 = x_2, \zeta x_1 = x_2$. So $\operatorname{Ker} f$ is $\{(0,0), (x_0, x_0), (2x_0, 2x_0)\}$. In same way for D, the action $\sigma \curvearrowright D \times D$ can be diagonalized. So we have

$$\Big((C \times C \times D \times D)/\langle \sigma, \tau \rangle\Big)/\langle \tau_2, \tau_3, \tau_4 \rangle = \Big((C \times C \times D \times D)/\langle \tau_3, \tau_4 \rangle\Big)/\langle \tau_2, \tilde{\sigma}, \tilde{\tau} \rangle.$$

Hence

$$\Big((C^2 \times D^2)/\mathfrak{S}_3\Big)\Big/\langle \tau_2, \tau_3, \tau_4 \rangle \cong (C \times C \times D \times D)/\langle \tilde{\sigma}, \tilde{\tau}, \tau_2 \rangle.$$

Since $\langle \tilde{\sigma}, \tau_2 \rangle$ is a normal subgroup of $\langle \tilde{\sigma}, \tilde{\tau}, \tau_2 \rangle$, we have

$$(C \times C \times D \times D)/\langle \tilde{\sigma}, \tilde{\tau}, \tau_2 \rangle \cong \Big((C \times C \times D \times D)/\langle \tilde{\sigma}, \tau_2 \rangle\Big)\Big/\langle \tilde{\tau} \rangle.$$

We can change generators of $\langle \tilde{\sigma}, \tau_2 \rangle$ from $\{\tilde{\sigma}, \tau_2\}$ to $\{\tau_2 \circ \tilde{\sigma}, \tau_2 \circ \tilde{\sigma}^2\}$. Each action of $\tau_2 \circ \tilde{\sigma}$ and $\tau_2 \circ \tilde{\sigma}^2$ are respectively given by

$$\begin{aligned} \tau_2 \circ \tilde{\sigma} \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (\zeta^2 x_1, x_2, y_1, \zeta y_2) \\ \tau_2 \circ \tilde{\sigma} \curvearrowright C \times C \times D \times D \ni (x_1, x_2, y_1, y_2) &\mapsto (x_1, \zeta x_2, \zeta^2 y_1, y_2). \end{aligned}$$

We identify $C \times C \times D \times D$ with $(C \times D) \times (C \times D)$ in the following way;

$$\begin{aligned} C \times C \times D \times D &\leftrightarrow (C \times D) \times (C \times D) \\ (x_1, x_2, y_1, y_2) &\leftrightarrow \big((x_1, y_2), (x_2, y_1)\big). \end{aligned}$$

The action of $\tilde{\tau}$ on $(C \times D) \times (C \times D)$ is given by

$$\tilde{\tau} \curvearrowright (C \times D) \times (C \times D) \ni (x_1, y_2, x_2, y_1) \mapsto (\zeta^2 x_2, \zeta y_1, \zeta x_1, \zeta^2 y_1).$$

So we have

$$\Big((C \times C \times D \times D)/\langle \tilde{\sigma}, \tau_2 \rangle\Big)\Big/\langle \tilde{\tau} \rangle \ \cong \ \operatorname{Sym}^2\Big((C \times D)/\mathbb{Z}_3\Big).$$

Here the $\mathbb{Z}_3$ action is the same one as in the first part of this section. So we get the conclusion. □

Reference

[Bea] Beauville, A. *Variétés kählerriennes dont la premiére classes de Chern est nulle*, J, of Diff. Geometry, **18**(1983), 755–782.

[B-D] Beauville, A. ,Donagi, R. *La variété des droites d'une hypersurface cubique de dimension 4*, C.R. Acad. Sc. Paris, **301**(1985), 703–706

[C-G] Clemens, C. Herbert. Griffiths, Phillip A. *The intermediate Jacobian of the cubic threefold*. Ann. of Math. (2), **95**(1972), 281–356.

[F-N] Fu,B., Namikawa,Y. *Uniqueness of crepant resolutions and symplectic singularities*. Ann. Inst. Fourier (Grenoble), **54**(2004), 1–19.

[Gri] P. Griffiths, *On the periods of certain rational integrals*, Ann. of Math. **90** (1969), 460–541.

[Har] Hartshorne, R. *Algebraic geometry*, Graduate Text of Mathematics (52) Springer.

[Huy] Huybrechts, D. *Compact Hyperkähler manifold*, Invent.Math., **135**(1999), 63–113.

[Kaw] Kawamata,Y. *On the length of an extremal rational curve*. Invent. Math., **105**(1991), 609–611.

[Kol] Kollár, J. *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihree Grezgebite, Band32, (1996), Springer

[Nak] Nakamura,I. *Hilbert schemes of abelian group orbits*, J. Algebraic Geom. **10**, (2001), 757–779.

[Nam] Namikawa,Y. *Deformation theory of singular symplectic n-folds*, Mathematische Annalen **319**(2001), 597–623.